22:Winterbotham et al. K-4 (no date) 1944 MS37 Senegal MS 39

Gambia, often in rotation with ground-nuts. Of the quick-maturing varieties *souna* and *sanio* are the most popular, the former thriving in sandy soil, the latter in heavier ground. Several kinds of sorghum are also cultivated. In the river area millet is sown where the floods have subsided and is harvested in April. In other parts it is planted immediately before the rainy season. Production probably averages about 300,000 tons annually. The export of millet from one province to another is controlled according to local needs.

Rice. The area flooded by the Casamance is naturally suited for rice cultivation. The rice-fields are handed down from father to son, each family growing sufficient for its own consumption and to pay its taxes. A store of rice is always kept to guard against a lean year or to be exchanged for some other commodity required.

Sugar-cane. This is grown near every village where there is enough water. No attempt has been made to commercialize the crop, and the West African varieties are more woody and contain less juice than those grown elsewhere for export. Sugar-cane (*Saccharum officinarum*) is a perennial plant of the grass family. It is easy to grow, even near the sea, but takes at least a year to mature. When grown for local use, as it is here, sugar is not extracted, but the red or white stems are chewed.

The Baobab. The baobab (*Adansonia digitata*) is found generally in the thornland (Plate 111), and is, therefore, much in evidence in Senegal. It is remarkable for the large girth of its trunk in comparison to its branches and leaves. The long green fruit, known as monkey-bread, is used as a food or as a seasoner. Boiled with water and then cooled, it also makes a refreshing drink, while the leaves and the young shoots are also good to eat. The inner bark yields a tough fibre, which can be twisted into rope or woven into a waterproof cloth. About 30 years ago a factory was opened at Dakar to prepare the fibre for export to rope- and paper-making firms in France. The bark and fibre can be stripped off without causing the death of the tree.

Manioc, beans, and maize are grown generally in the vicinity of villages. In the Cape Verde peninsula these, as well as yams, sweet potatoes, and melons, are grown for the Dakar markets.

Economic Crops

Ground-nuts. Ground-nuts (*Arachis hypogaea*) require a tropical climate and an annual rainfall of from 14 to 60 inches. They are the

22:Winterbotham et al. K-4 (no date) 1944 MS37 Senegal MS 37

sixth crop of Africa, and 43 per cent. of the total continental production comes from Senegal. The two main groups cultivated are the 'runner' and the 'erect'. The Rufisque and Cayor varieties, both runners, are those grown generally here and also in British West Africa. The method of cultivation is described in Volume I, p. 321.

Ground-nuts were originally grown as a staple native crop, but the industrial demand for vegetable oils, coupled with the development of transport facilities, opened up a world market for them. Extra labour for the harvest is provided annually by seasonal immigration from French Sudan, French Guinea, Portuguese Guinea, and the Gambia. In 1938 the number of outside workers was 69,757. Of these, 56,743 went to the Kaolack area, between 1,000 and 3,500 to Sedhiou, Ziguinchor, Kolda, Tambacounda, and Diourbel, and the remainder to Thiès and Linguéré.

There are numerous shelling factories. Early in 1939 new ones were opened at Kaolack, Ziguinchor, and Koungheul.

In 1938 nearly 2,000,000 acres were under cultivation and the experimental station at Bambey distributed 5,414 tons of selected seed. In 1939 7,382 tons of seed were sent out, with a corresponding increase in acreage.

During the British blockade of French West Africa after the fall of France in 1940, ground-nuts were converted into fuel oil. If this practice has been proved worth continuing after the war, cultivation may be greatly extended.

Sisal. Sisal originated in Mexico, being named after the port of Sisal in Yucatan province, and it was introduced into Senegal at the end of the nineteenth century. The species grown are *Agave rigida sisalana*, *A. rigida elongata*, and *A. rigida zapupe*. The rigid leaves containing the fibre are about 4 feet long and are harvested biennially from the fourth to the eighth or tenth year. When a plant has finished producing it shoots up a tall flower stem, with bulbils or buds on the end, and then withers.

The preparation of stem and leaf fibres is lengthy. First the fleshy tissue encasing the fibre is scraped away, and then the fibre is washed, hung out to dry, and brushed. It is now ready for grading. Next it is combed into a continuous sliver of untwisted rope. The final process is the drawing and twisting of the sliver to a uniform thinness for spinning into a rope that is particularly resistant to sea-water. The largest plantation is in French Sudan, and stretches for 12 miles along the bank of the Senegal near Kayes. Other smaller ones are in Tambacounda and on the Cape Verde peninsula. There is oppor-